

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A01N 63/02, C12R 1/01, 1/38, 1/385, 1/39 // (C12R 1/01, 1:38, 1:385, 1:39)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/22598</b> <b>(43) International Publication Date:</b> 14 May 1999 (14.05.99)
<b>(21) International Application Number:</b> PCT/GB98/03254 <b>(22) International Filing Date:</b> 30 October 1998 (30.10.98)  <b>(30) Priority Data:</b> 9722960.3 30 October 1997 (30.10.97) GB 9723226.8 3 November 1997 (03.11.97) GB  <b>(71) Applicant (for all designated States except US):</b> THE UNIVERSITY OF READING [GB/GB]; Whiteknights, P.O. Box 217, Reading RG6 6AH (GB).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> HAGUE, Nigel, Graham, Mackenzie [GB/GB]; 19 Greenleas Avenue, Emmer Green, Reading, Berkshire RG4 8TA (GB). ELAWAD, Sami, Abdulrahaman [SD/GB]; Sibly Hall, Redhatch Drive, Reading, Berkshire RG2 5QW (GB).  <b>(74) Agents:</b> O'BRIEN, Caroline, J. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> BIOPESTICIDE: MATERIALS AND METHODS		
<b>(57) Abstract</b>		
<p>The invention provides biopesticides for the control of insect pests and/or plant parasitic nematodes. The effective agent is a motile species of bacteria or a bacterial symbiont of an entomopathogenic nematode.</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

BIOPESTICIDE: MATERIALS AND METHODS

The present invention concerns materials and methods relating to biopesticides. In particular, the present invention concerns materials and methods relating to  
5 biopesticides comprising bacteria.

The damage done to plants and humans by harmful insects and/or plant parasitic nematodes is enormous and their control is mainly achieved through use of synthetic chemical pesticides.

10 However there are many problems related to the use of chemical pesticides including resistance and toxicity to non-target species. These and other problems and issues create a political and social climate opposed to them. Therefore microbial alternatives including  
15 entomopathogenic nematodes have been sought. These nematodes, of the genera *Steinernema* and *Heterorhabditis*, are widespread and have been isolated on every inhabited continent and on many islands. Commercialisation of such nematodes are exempt from registration requirements in  
20 many countries. Efficacy has been demonstrated in various markets, so product and sales have begun on a commercial scale.

*Steinernematid* and *heterorhabditid* nematodes are symbiotically associated with bacteria in the genera  
25 *Xenorhabdus* and *Photorhabdus* respectively. The free-living, infective third stage Dauer juvenile of the nematode, which does not feed, carries the bacterium

within its intestine. The infective-stage locates insects, initiates infection and is the only stage in the nematode's life cycle that survives outside the insect in the soil. The infective-stage enters insects via mouth, anus or spiracles and penetrates mechanically into the body cavity where it releases the bacterium. In certain insects, *heterorhabditids* can enter into the insect body cavity through the cuticle. The bacterium proliferates, causes septicemic death of the insect within 24-72 hours and establishes favourable conditions for nematode reproduction by providing nutrients and inhibiting the growth of many foreign micro-organisms. The nematodes feed on multiplying bacteria and dead host tissue, passing through several generations. Eventually, Dauer Juveniles (DJs), carrying symbiont bacteria in their gut, emerge from the insect cadaver. At 18-28°C, the nematodes complete their life cycle in most insects in 8-20 days.

Exposure of insect hosts to high concentrations of *steinernematids* and *heterorhabditids* under laboratory conditions has indicated that the host range includes most insects and at high concentrations extends also to other invertebrates (*Gastropoda*, *Symphyla*, *Arachnida*, *Crustacea* and *Diplopoda*). However, behavioural and ecological barriers have restricted the effectiveness of nematodes to certain soil inhabiting insects. Consequently, laboratory experimental infections do not always appear to translate to the field where the nematodes infect fewer species.

In order to achieve field results comparable to that of standard chemical pesticides the strategy for application of entomopathogenic nematodes has to be very carefully controlled.

5 Further the current cost of preparing entomopathogenic nematodes for pest control is 10-60% higher than chemical insecticides and some nematodes of interest e.g. heterorhabditid nematodes cannot yet be produced efficiently in liquid fermenters.

10 Microbial pesticides other than entomopathogenic nematodes have been used and Table 1 summarises the major commercially available microbial insecticides (from Lacey, L.A. and Goettel M.S., 1995 Entomophaga 40, p3-27 and Georgis R 1997 BCPC Symposium Proceedings No 68 p  
15 243-252).

Of the microbes referred to in Table 1, the bacteria *Bacillus thuringiensis* (Bt) is most widely used. Various subspecies which are larvicidal for different host ranges have been introduced. The effective Bt larvicidal agent  
20 comprises a toxin and Bt genes that encode toxin production have been manipulated using both recombinant and non-recombinant methods. Manipulation of these Bt toxin genes has enabled their incorporation into crop plants.

25 However in the case of Bt-transgenic plants, constant control of insect pests will be an extremely strong selective force for insects that are resistant to Bt, necessitating a need for aggressive resistance

management plans where growers set aside acres that are either not treated with insecticides or not treated with Bt.

5 Owing to the problems of development of resistance there is a real economic need for as diverse a range of biopesticides as possible.

The present inventors have now unexpectedly discovered that isolates of the symbiotic bacteria can be used to control insect pests and/or plant parasitic  
10 nematodes.

Thus the present invention provides a new biopesticide composition for the control of insect pests and/or plant parasitic nematodes which comprises as an effective agent a bacterial species which is motile.

15 The present invention provides a new biopesticide composition which comprises as an effective agent a bacterial species which is flagellate.

The present invention provides a new biopesticide composition which comprises as an effective agent a  
20 bacterial species which enters the target insect species via a route other than the alimentary canal. It may enter via the spiracles.

The present invention provides a new insecticide composition which comprises as an effective agent a  
25 bacterial symbiont of an entomopathogenic nematode. The bacterial symbiont may be motile. It may be flagellate. It may enter a target insect species via a route other than the alimentary canal. It may enter via the

spiracles.

The present invention provides a new biopesticide which comprises a bacterial species which produces one or more toxins which interfere with the behaviour and activity of a plant parasitic nematode.

The bacterial species may be selected from the genera *Xenorhabdus*, *Photorhabdus*, *Pseudomonas* or *Flavimonas*.

The bacterial species may be of the genus *Xenorhabdus* or the genus *Flavimonas*. The bacterial species may be of the genus *Flavimonas*.

The genus *Flavimonas* contains only one species to date *F. oryzihabitans*. The genus was split-off "taxonomically" from the genus *Pseudomonas*. Thus the two genera *Flavimonas* and *Pseudomonas* are closely related. *Pseudomonas* has been developed as a biological agent to control fungi. However *Pseudomonas* spp. have not been suggested as agents to control insects and/or plant parasitic nematodes. In relation to use as an agent to control insect pests and/or plant parasitic nematodes, bacteria of the genus *Flavimonas* may have some advantages over bacteria of the genus *Xenorhabdus*. Firstly *Flavimonas* appears to be more active, with better stability in water and resistance to drying.

The bacterial species may be *Xenorhabdus nematophilus* or *Flavimonas oryzihabitans* or *Xenorhabdus bovienii*. The bacterial species may be *Flavimonas oryzihabitans*.

The biopesticide composition as stated above may comprise a plurality of different bacterial species thus a biopesticide composition according to the invention may comprise a plurality of species selected from bacteria of the genera *Xenorhabdus*, *Photorhabdus*, *Pseudomonas*, *Flavimonas*. Thus the biopesticide composition may comprise a plurality of species selected from *Xenorhabdus nematophilus*, *Flavimonas oryzihabitans* or *Xenorhabdus bovienii*.

The biopesticide composition may be adapted for application as a liquid suspension of bacteria. The biopesticide composition may be adapted for application as an aqueous suspension of bacteria.

The present invention also provides the use of a bacterial species as stated above (i.e. motile bacteria, flagellate bacterial, spiracle-infecting bacteria, selected from the genera *Xenorhabdus*, *Pseudomonas*, *Photorhabdus* or *Flavimonas*) in the preparation of a biopesticide composition wherein the bacterial species comprises a biopesticidally effective element of the composition.

The invention also provides a method for the manufacture of a biopesticide composition, characterised in the use, as an essential constituent of said composition of a bacterial species as stated above (i.e. motile bacteria, flagellate bacterial, spiracle-infecting bacteria, selected from the genera *Xenorhabdus*, *Pseudomonas*, *Photorhabdus* or *Flavimonas*) in the



preparation of a biopesticide composition wherein the bacterial species comprises a biopesticidally effective element of the composition.

The present invention also provides a method of  
5 controlling insect pests and/or plant parasitic nematodes which comprises applying to environment of the pest or parasite, a biopesticide composition as stated above. The method may comprise application of the composition to a crop plant. In particular the biopesticide composition  
10 for application may comprise a liquid suspension of bacteria, for example an aqueous suspension.

The suspension may contain about  $3-8 \times 10^7$  bacterial cells per ml. The suspension may contain about  $5-7 \times 10^7$  bacterial cells/ml. The suspension may contain about  $4 \times$   
15  $10^7$  bacterial cells per ml.

However a skilled person will be readily able to ascertain the most advantageous dosage rate for a given bacteria or combination of bacteria in relation to the control of a particular pest or combination of pests  
20 using the teaching of the specification and common general knowledge.

#### Example 1

Bacteria suitable for use in relation to the present invention can be obtained from commercial sources and  
25 depository institutions.

Alternatively they can be isolated from biological sources such as entomopathogenic nematodes. Such nematodes are available from commercial sources and depository institutions. They can also be readily  
5 isolated from insect and soil samples.

Suitable bacteria occur in the infective juveniles of the nematodes and in soils throughout the world. They can be isolated from soil by using suitable hosts, such as the wax moth, *Galleria mellonella*, as trap insects.

10 1. *Xenorhabdus bovienii*

This bacteria species is associated with the nematode *Steinernema feltiae* and the bacterium can be obtained from MicroBio, Unit 2 Centro, Boundary Way, Hemel Hempstead, Hertfordshire HP2 7SU, who  
15 produce *S. feltiae* as the product NEMASYS.

2. *Xenorhabdus nematophilus*

This bacterial species is associated with the nematode *Steinernema capocapsae* and can be obtained from Thermotrilogy, 7500 Grace Drive, Columbia MD  
20 21044-4098, USA. Thermotrilogy can also supply bacteria from another Steinernematid, *S. riobrave*.

3. *Photorhabdus luminescens*

This bacterial species can be obtained from MicroBio (see above) who produce the nematode *Heterorhabditis megidis* as NEMASYS H.  
25

4. *Flavimonas oryzihabitans*

The genus *Flavimonas* was first described in a paper by Holmes, B et al (1987) in International Journal of Systematic Bacteriology Vol 37 No 3, p245-250

5 ("Chryseomonas luteola comb. nov. and *Flavimonas oryzihabitans* gen. nov., comb. nov., Pseudomonas-Like Species from Human Clinical Specimens and Formerly Known, respectively as Groups Ve-1 and Ve-2") Strains of *F. oryzihabitans* have been deposited

10 in the National Collection of Type Cultures: L327/81 as NCTC 11851 and E5726 as NCTC 11852. B2524 is available from the American Type Culture Collection as NTCC 35564.

In an earlier reference the bacteria now designated

15 *F. oryzihabitans* was described as *Pseudomonas oryzihabitans* and obtainable from rice paddy field soils and clinical specimens (Kodma K., et al (1985) International Journal of Systematic Bacteriology Vol 35 No 4: "Two New Species of

20 *Pseudomonas*: *F. oryzihabitans* Isolated from Rice Paddy and Clinical Specimens and *F. luteola* Isolated from Clinical Specimens").

More recently *F. oryzihabitans* has been isolated from the *Steinernematid*, *S. abbasi*. This is a new

25 association between a bacterium and an entomopathogenic nematode: Elawad, S., et al in Fundam. Appl. Nematol. (1997) 20 p435-442 "*Steinernema abbasi* sp.n. Nematode:

Steinernematidae from The Sultanate of Oman)" provides details concerning the isolation and characteristics of *S.abbasi* which comprise a source of *Flavimonas oryzihabitans*.

5           The following details concern the preparation of a bacterial isolate from nematodes carrying the appropriate bacterial species. Detailed methodologies of relevance are in: "Steinernematid and Heterorhabditid Nematodes: A Handbook of Biology and Techniques by Jennifer L.  
10   Woodring and Harry K. Kaya, Southern Cooperative Series Bulletin 331 published August 1988 by the Nematode Subcommittee of the Southern Regional Project S-135 Entomopathogens for Use in Pest-Management Systems, Arkansas Agricultural Experiment Station, Fayetteville,  
15   Arkansas.

Bacteria are isolated from the haemocoel of a larva of the greater wax moth, *Galleria mellonella* infected with the bacterium of interest. Briefly the method is as set out below.

- 20   1.   Late instar larvae of *G.mellonella* are infected with the appropriate Dauer juveniles (DJs) at the rate of approximately 200 DJs per petri-dish lined with a double layer of filter paper.
2.   After the death of the larvae they are washed and  
25   surface-sterilized with about 70% alcohol in

staining blocks for 5-10 mins.

3. Sterilized larvae are left to dry in a laminar flow cabinet for about 1 min in a sterile petri-dish.
4. The dead larva are opened with sterile scissors and  
5 needle in the laminar flow cabinet and then left for about ½ minute.
5. A drop from the oozing haemolymph from the cadaver is streaked with a sterile toothpick onto a  
10 previously prepared nutrient both agar (NBTA) made up as follows: 37 g standard agar, 25 mg bromothymol blue, 1000 ml distilled water, 4 ml of 1% filtered 2,3,5 triphenyl-tetrazolium chloride solution.
6. The agar plates are sealed with parafilm and incubated at 28°C in the dark for 24 hr.
- 15 7. Single colonies are selected from the culture to grow in NBTA and then used to streak a new plate of NBTA. The reculturing is done to obtain a culture of the bacterium which was uniform in colony morphology.
- 20 8. From the pure cultures of the bacterium a single colony is selected and picked out by sterile toothpick and used to inoculate 50 ml of liquid

12

broth held in glass flasks. The flasks are covered with sterile cotton wool and paper held by a rubber band. The flasks are placed in a shaking incubator for 2 days at 28°C; the shaking being adjusted to 150 rpm.

5

The concentration of the bacteria in the broth is estimated using a haemocytometer. Extra sterile broth can be used to dilute the concentration of the bacteria in the solution to give the required dose for application.

10

#### Example 2

Bacterial suspensions of *Flavimonas oryzihabitans* and *Xenorhabdus nematophilus* were used to investigate control of *Spodoptera exigua* (cutworm).

15

#### Investigation A

20 leaves were detached from the cotton plant *Gossypium hirsutum* and put in separate small covered plastic containers containing sterile distilled water so that the leaves remained fresh during the experiment. The small container with the leaf was then put in a larger plastic container and sealed.

20

Both the top and bottom surfaces of the leaf were sprayed with 1 ml of bacterial suspension comprising  $4 \times 10^7$  bacterial cells per ml of solution (in 3% (V/V) Tween 80

as emulsifier). Each cotton leaf had a surface area of approximately 59 cm<sup>2</sup> and one 3rd instar larva of *S.exigua* was placed on each leaf after spraying with the bacterial suspension.

5 The experiment was placed in a controlled temperature room at 28°C.

The control treatment received Tween 80 in distilled water.

10 Assessment of the number of dead *S.exigua* larvae was done after 72 hours.

The experiment was conducted using (a) *F. oryzihabitans* and (b) *X. nematophilus*.

#### Investigation B

15 This experiment was conducted as in Investigation A above except that, after application of the bacterial suspensions, the leaves were all allowed to dry in a laminar flow cabinet prior to the introduction of one *S. exigua* larva per cotton leaf. As in Investigation A the number of dead *S. exigua* larvae was derived after 72  
20 hours.

#### Investigation C

This experiment was conducted to investigate the effect of the bacteria on *S.exigua* pupae.

25 55 g of sterile sand was mixed with 15 ml of bacterial suspension comprising  $4 \times 10^7$  bacterial cells per ml of solution (in 3% (V/V) Tween 80) and placed in a petri

dish. 10 pupae of *S. exigua* were buried in the moist sand and the emergence of *S. exigua* adults assessed after 5 days. 4 such experimental dishes were established.

The results are summarised in Table 2.

5           In moist conditions *F. oryzihabitans* and *X. nematophilus* caused 100% mortality within 72 hours. Mortality of 75% was found in about 48 hours (data not shown). Application of bacteria to sand in which pupae of *S. Exigua* are placed resulted in mortalities in excess  
10       of 75%.

The present inventors have also found that if *F. oryzihabitans* or *X. nematophilus* are incorporated in the artificial diet used to rear *S. exigua*, mortalities in excess of 75% resulted (data now shown).

15           Although both *F. oryzihabitans* and *X. nematophilus* can be kept fully viable in nutrient broths, they eventually lose their viability in sand (*X. nematophilus* loses viability more rapidly than *F. oryzihabitans*).

### Example 3

20           Bacterial solution of *Flavimonas oryzihabitans* and *Xenorhabdus nematophilus* were used to control *Pieris brassicae* (large white butterfly).

The experiment was conducted using larvae as set out in Example 2, Investigation A above).

25           The results are summarised in Table 3.



Example 4

Bacterial solution of *Flavimonas oryzihabitans* and *Xenorhabdus nematophilus* were used to control *Galleria mellonella* (greater wax moth).

5        The experiment was conducted using larvae substantially as set out in Example 2, Investigation C above except that 40 test larvae were used and mortality assessment was carried out after 10 days.

The results are summarised in Table 4.

10        The experimental results given in Table 4 show that when these bacteria are applied to insect larvae and pupae under conditions when the host larvae, pupae and bacteria are kept moist, there is a resultant high insect mortality.

15        Example 5

Three experiments were conducted to investigate the effect of symbiotic bacteria on juveniles of plant parasitic nematodes.

20        Bacterial suspensions of *Flavimonas oryzihabitans* and *Xenorhabdus nematophilus* were mixed with nutrient agar to obtain a concentration of  $1.6 \times 10^{10}$  cells/ml. Second stage juveniles ( $J_2$ s) of *Meloidogyne javanica* (root-knot nematode) and *Globodera rotochiensis* (potato cyst nematode) were placed on the agar and their movement  
25        observed at 30 min; 1h; 2h; 4h; 8h; 16h and 24h. There was no movement of juveniles of either species after 30

min exposure.

In a second experiment, juveniles of *M.javanica* were exposed to concentrations of  $1.6 \times 10^8$  and  $1.6 \times 10^{10}$  cells per ml of *F.oryzihabitans* and a concentration of  $1.6 \times 10^{10}$  of *X.nematophilus*. After exposure for 1h, 3h, 6h, 12h, 24h and 7 days the nematode J<sub>2</sub>s were transferred to water to observe whether they recovered some movement. Observation for recovery of movement was carried out at 1h, 3h, 6h, 24h and 7 days post transfer to water.

After treatment with *F.oryzihabitans* for 1h and 3h the nematodes showed partial recovery of movement and partial recovery of movement was observed after 1h treatment with *X.nematophilus*. This was observed for both concentrations of *F.oryzihabitans*. Where (a) juveniles were exposed to  $1.6 \times 10^8$  cells/ml *F.oryzihabitans* for 6 hours and greater; (b) juveniles were exposed to  $1.6 \times 10^{10}$  cells/ml *F.oryzihabitans* for 6 hours and greater; and (c) juveniles were exposed to  $1.6 \times 10^{10}$  *X.nematophilus* for 3 hours and greater there was no recovery of movement upon transfer to water.

In a third experiment bacterial suspensions of *F.oryzihabitans* ( $1.6 \times 10^{10}$  cells/ml) were stored at 7°C for 10 days and then tested against J<sub>2</sub>s of *M.javanica*. There was no movement of J<sub>2</sub>s after a minimum exposure of 1h.

Table 1  
Major commercially available microbial insecticides

Pathogens	Major targeted group
Bacteria	
<i>Bacillus thuringiensis</i>	
Crystal Protein types:	
Cry IA-G, Cry IIA-C	Lepidoptera
Cry IIIA-D	Coleoptera, Culicidae, Simuliidae
Cry IVA-D, Cry IIA	
<i>B. sphaericus</i>	Culicidae
Virus	
Nuclear polyhedrosis viruses	Lepidoptera, Hymenoptera
Granulosis viruses	Lepidoptera
Fungi	
<i>Beauveria bassiana</i>	Coleoptera, Lepidoptera, Homoptera, Orthoptera
<i>Metarhizium anisopliae</i>	Coleoptera, Lepidoptera, Homoptera, Orthoptera
<i>Verticillium lecanii</i>	Homoptera, thrips
Nematodes	
<i>Steinernema carpocapsae</i>	Lepidoptera, Coleoptera, (Curculionidae, Chrysomelidae)
<i>S. feltiae</i>	Diptera (Sciaridae)
<i>S. riobravus</i>	Lepidoptera, Orthoptera (mole crickets), Coleoptera (Curculionidae)
<i>Heterorhabditis bacteriophora</i>	Lepidoptera, Coleoptera
<i>H. megidis</i>	Coleoptera

Table 2

The mortality of *S. exigua* larvae and pupae after treatment with the bacteria *F. oryzihabitans* and *X. nematophilus*.

Treatment	Mortality	
	Application on leaves which were kept moist	Pupae Application followed by drying
<i>Flavimonas oryzihabitans</i>	100%	20% 80%
<i>Xenorhabdus nematophilus</i>	100%	10% 80%
Untreated	0%	0% 10%

Table 3

The mortality of *Pieris brassicae* larvae after treatment with the bacteria  
*F. oryzihabitans* or *X. nematophilus*.

<u>Treatment</u>	<u>Mortality</u>
Flavimonas oryzihabitans	90%
Xenorhabdus nematophilus	70%
Untreated	0%

Table 4

The mortality of *Galleria mellonella* larvae after treatment with the bacteria  
*F. oryzihabitans* or *X. nematophilus*.

<u>Treatment</u>	<u>Mortality</u>
Flavimonas oryzihabitans	70%
Xenorhabdus nematophilus	60%
Untreated	0%

CLAIMS:

1. A biopesticide for the control of insect pests or  
plant parasitic nematodes or both, which comprises as an  
effective agent a species of bacteria which is a symbiont  
5 of an entomopathogenic nematode.
2. A biopesticide according to claim 1 wherein the  
bacteria is a motile species.
3. A biopesticide according to claim 1 or claim 2  
wherein the bacterial species is of a genera selected  
10 from *Xenorhabdus*, *Photorhabdus*, *Pseudomonas* or  
*Flavimonas*.
4. A biopesticide according to any one of claims 1 to 3  
which is an insecticide.
5. A biopesticide according to any one of claims 1 to 3  
15 which is a vermicide.
6. A biopesticide according to any one of claims 1 to 5  
wherein the bacteria is flagellate.
7. A biopesticide according to any one of claims 1 to 6  
wherein the bacteria invades insects via a route other  
20 than the alimentary canal.

8. A biopesticide according to claim 7 wherein the bacteria invades insects via the spiracles.
9. A biopesticide according to any one of claims 1 to 8 wherein the bacteria produces a toxin which interferes with the behaviour and activity of a plant parasitic nematode.
10. A biopesticide according to any one of claims 1 to 9 wherein the bacterial species is of the genus *Xenorhabdus*.
11. A biopesticide according to any one of claims 1 to 9 wherein the bacterial species is of the genus *Flavimonas*.
12. A biopesticide according to claim 11 which comprises *Flavimonas oryzihabitans*.
13. A biopesticide according to claim 10 which comprises *Xenorhabdus nematophilus*.
14. A biopesticide according to claim 10 which comprises *Xenorhabdus bovienii*.
15. A biopesticide according to any one of claims 1 to 14 which comprises as an effective agent a mixture of bacteria.



16. A biopesticide according to claim 15 which comprises a plurality of bacterial species of genera selected from *Xenorhabdus*, *Photorhabdus*, *Pseudomonas*, *Flavimonas*.

5 17. A biopesticide according to any one of claims 1 to 16 adapted for application as a liquid suspension of bacteria.

18. A biopesticide according to any one of claims 1 to 17 adapted for application as an aqueous suspension of bacteria.

10 19. Use of a species of bacteria which is a symbiont of an entomopathogenic nematode in the preparation of a biopesticide for the control of insect pests and/or plant parasitic nematodes.

15 20. Use according to claim 19 wherein the bacteria is motile species.

21. Use according to claim 19 wherein the bacterial species is of a genera selected from *Xenorhabdus*, *Photorhabdus*, *Pseudomonas* or *Flavimonas*.

20 22. A method of controlling insect pests or plant parasitic nematodes or both which comprises applying to the environment of the pest or parasite a biopesticide according to any one of claims 1 to 21.

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/GB 98/03254

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N63/02 C12R1/01 C12R1/38 C12R1/385 C12R1/39  
 //(C12R1/01, 1:38, 1:385, 1:39)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N C12R

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 98 08388 A (MORGAN JAMES ALUN WYNNE ;JARRETT PAUL (GB); ELLIS DEBORAH JUNE (GB) 5 March 1998 see examples 1,3,6 ---	1-10, 13-23
X	US 5 616 318 A (DUDNEY RALPH A) 1 April 1997 see the whole document ---	1-10, 13-23
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

9 February 1999

Date of mailing of the international search report

24/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
 Fax: (+31-70) 340-3016

Authorized officer

Klaver, J

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 98/03254

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	R.-U. EHLERS, A. WULFF & A. PETERS: "Pathogenicity of axenic <i>Steinernema feltiae</i> , <i>Xenorhabdus bovienii</i> , and the bacto-helminthic complex to larvae of <i>Tipula oleracea</i> (Diptera) and <i>Galleria mellonella</i> (Lepidoptera)." J. INVERTEBRATE PATHOL., vol. 69, 1997, pages 212-217, XP002092716 see: page 214, col. 2, par. 1 - 3 and Figure 1; page 215, col. 2, par. 3 - page 216, col. 1, par. 1. ---	1-10, 13-23
X	M. TACHIBANA, H. HORI, N. SUZUKI, T. UECHI, D. KOBAYASHI, H. IWAHANA & H. K. KAYA: "Larvicidal activity of the symbiotic bacterium <i>Xenorhabdus japonicu</i> from the entomopathogenic nematode <i>Steinernema kushidai</i> against <i>Anomala</i> <i>cuprea</i> (Coleoptera: Scarabaeidae)." J. INVERTEBRATE PATHOL., vol. 68, 1996, pages 152-159, XP002092717 see: page 153, col. 2, par. 5 - page 154, col. 1, par. 1; page 154, col. 1, par. 1 - col. 2, par. 1; page 157, col. 2, par. 3 ---	1-10, 13-23
X	CHEMICAL ABSTRACTS, vol. 118, no. 1, 4 January 1993 Columbus, Ohio, US; abstract no. 3550, S. YAMANAKA ET AL.: "Biochemical and physiological characteristics of <i>Xenorhabdus</i> species, symbiotically associated with entomopathogenic nematodes including <i>Steinernema kushidai</i> and their pathogenicity against <i>Spodoptera litura</i> (Lepidoptera: Noctuidae)" XP002048914 see abstract	1-10, 13-23
X	& ARCH. MICROBIOL., vol. 158, no. 6, 1992, pages 387-393, ---	1-10, 13-23
X	R. J. AKHURST & N. E. BOEMARE: "Biology and Taxonomy of <i>Xenorhabdus</i> " 1990, CRC PRESS, INC., BOCA RATON XP002092721 in: R. Gaugler & H. K. Kaya (eds.): "Entomopathogenic Nematodes in Biological Control", pages 75 - 89. See page 75, par. 1 - page 77, par. 3. ---	1-10, 13-23
	---	
	-/--	

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/GB 98/03254

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S. FORST & K. NEALSON: "Molecular biology of the symbiotic-pathogenic bacteria <i>Xenorhabdus</i> spp. and <i>Photorhabdus</i> spp." MICROBIOLOGICAL REVIEWS, vol. 60, no. 1, March 1996, pages 21-43, XP002092718 see page 26, column 1, paragraph 1 - page 28, column 2, paragraph 4 ----	1-10, 13-23
X	N. BOEMARE, C. LAUMOND & H. MAULEON: "The entomopathogenic nematode-bacterium complex: biology, life cycle and vertebrate safety." BIOCONTROL SCI. AND TECHNOL., no. 6, 1996, pages 333-345, XP002092719 see page 336, paragraph 2 - page 339, paragraph 2 ----	1-10, 13-23
Y	Y. ANZAI, Y. KUDO & H. OYAIZU: "The phylogeny of the genera <i>Chryseomonas</i> , <i>Flavimonas</i> and <i>Pseudomonas</i> support synonymy of these three genera." INT. J. SYST. BACTERIOL., vol. 47, no. 2, April 1997, pages 249-251, XP002092720 see the whole document -----	11,12
Y		11,12

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03254

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9808388	A	05-03-1998	AU 4024997 A	19-03-1998
US 5616318	A	01-04-1997	NONE	

**THIS PAGE BLANK (USPTO)**